

ADVANCED SUBSIDIARY GCE CHEMISTRY

How Far, How Fast?

THURSDAY 11 JANUARY 2007

2813/01

Morning

Time: 45 minutes

Additional materials: Scientific calculator

Data Sheet for Chemistry (Inserted)



Candidate Name	1 10 20 1 100	Continuos et Guino 445	1.476
Centre Number		Candidate Number	

INSTRUCTIONS TO CANDIDATES

- Write your name, Centre number and Candidate number in the boxes above.
- Answer all the questions.
- Use blue or black ink. Pencil may be used for graphs and diagrams only.
- Read each question carefully and make sure you know what you have to do before starting your answer.
- Do not write in the bar code.
- Do not write outside the box bordering each page.
- WRITE YOUR ANSWER TO EACH QUESTION IN THE SPACE PROVIDED. ANSWERS WRITTEN ELSEWHERE WILL NOT BE MARKED.

INFORMATION FOR CANDIDATES

- The number of marks for each question is given in brackets [] at the end of each question or part question.
- You may use a scientific calculator.
- A copy of the Data Sheet for Chemistry is provided as an insert with this
 question paper.
- You are advised to show all the steps in any calculations.

Qu.	Max	Mark
1	10	
2	14	
3	10	
4	11	
TOTAL	45	

This document consists of 8 printed pages and a Data Sheet for Chemistry.

Answer all the questions.


- 1 A group of students investigated the effect of concentration on the rate of a reaction. They used the reaction between magnesium carbonate and dilute hydrochloric acid and measured the rate at which the gas was collected.
 - (a) (i) An incomplete equation for this reaction is given below. Complete the equation by balancing it and inserting state symbols.

$$MgCO_3(....)$$
 + $HCl(....)$ \rightarrow $MgCl_2(....)$ + $CO_2(....)$ + $H_2O(....)$ [2]

(ii) Write the ionic equation for this reaction.

[1]

(b) The students added dilute hydrochloric acid to some magnesium carbonate. The students collected the gas and measured the volume, at regular intervals, until after the reaction was complete. They then plotted a graph of their results.



Use collision theory to explain the changes in the rate of the reaction as it proceeds.


(c)	The	The students repeated the experiment using a weak acid instead of hydrochloric acid.					
	Ass	sume the concentration of both acids and all other conditions are the same.					
	(i)	On the axes in (b), sketch the graph the students obtained from this experiment.	[2]				
	(ii)	State and explain what effect changing the acid has on the rate of the reaction.					
			•••••				
			[2]				
		По	tal: 101				

- 2 Propane, C₃H₈, is a gas at room temperature and pressure. It is used in blow torches to melt the bitumen needed to apply the felt to flat roofs.
 - (a) Write the equation for the complete combustion of propane.

(b) Define the term standard enthalpy change of combustion.

.....[3]

(c) A blow torch was used to determine the enthalpy change of combustion of propane. The apparatus is shown below.



In the experiment, 200 g of water were used. The temperature of the water changed from 18.0 °C to 68.3 °C when 1.00 g of propane was burnt.

(i) Calculate the energy produced in kJ. The specific heat capacity of water is 4.18 J g⁻¹ K⁻¹.

energy = kJ [2]

	(ii)	Calculat	e the numbe	r of moles of C ₃ H ₈ burnt during the experiment.
				mol [1]
	(iii)	Deduce	the enthalpy	change of combustion, in kJ mol ⁻¹ , of C ₃ H ₈ .
				enthalpy change of combustion = kJ mol ⁻¹ [2]
d)		ues of en	thalpy chang	ges of combustion can be used to calculate enthalpy changes of
		e enthalpy pane.	change for	the reaction in equation 2.1 is the enthalpy change of formation of
			3C(s) +	$4H_2(g) \longrightarrow C_3H_8(g)$ equation 2.1
		e table be pane.	elow shows	the enthalpy changes of combustion of carbon, hydrogen and
				enthalpy change of combustion / kJ mol ⁻¹
			carbon	-394
			hydrogen	-286
			propane	-2219
	(i)	Use the	se data to ca	Iculate the enthalpy change of formation of propane.
				enthalpy change of formation = kJ mol ⁻¹ [3]
	(ii)	Suggest		halpy change of formation of propane cannot be measured directly.
				[1]

[Turn over

[Total: 14]

		re used as flavourings. They are made by a reversible reaction between a carboxylic acid lcohol.
		carboxylic acid + alcohol
(a)	Give	e two features of a reversible reaction, when a dynamic equilibrium has been set up.
	1	
	2	
		[2]
(b)	The	production of esters is catalysed homogeneously by the presence of acids.
	(i)	What is meant by a catalyst?
		[1]
	(ii)	What is meant by homogeneous?
		[1]
	(iii)	Using the fact that acids are needed to catalyse this reaction, deduce the formula of the ion that acts as the catalyst.
		[1]
	(iv)	Catalysts do not affect the position of an equilibrium. Explain why not.
		[1]

Use the graph to	explain the effec	t of a cata	lyst on the ra	te of a reacti	on.

OCR OCR OCR OCR OCR

OCR OCR OCR OCR

Am	monia, NH ₃ , is made industrially by the Haber process. This is an equilibrium reaction.
	$N_2(g) + 3H_2(g) \implies 2NH_3(g)$ $\Delta H = -92 \text{ kJ mol}^{-1}$
(a)	State the pressure and temperature that are used in the Haber process.
	pressure
	temperature[2]
(b)	Describe and explain why these conditions are a compromise between rate and equilibrium.
	[9]
	[Total: 11]

END OF QUESTION PAPER

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

OCR is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.